No.



9500095

<u> AHUR UNIHURU SHAMUS ODRAMURRIOA</u>

TO ALL TO WHOM THESE; PRESENTS SHALL COME;

Aestern Plant Breeders, Inc.

Decemb, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED, PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR MPORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT D BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

WHEAT

'WestBred 936'

In Testimonn Mucrost, I have hereunto set my hand and caused the seal of the Mant Pariety Arotection Office to be affixed at the City of Washington, D.C. this thirty-first day of August in the year of our Lord one thousand nine hundred and ninety-five.

Allest:

RAAHQ

Acting Commissioner
Plant Variety Protection Offi

Secretary of Syriculture

U.S. DEPARTMENT AGRICULTURAL MA SCIENCE	Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C.		
APPLICATION FOR PLANT VAR	NS ON REVERSE)	TION CERTIFICATE	2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).
1. NAME OF APPLICANT(S) (as it is to appear on the Certificate		Z. TEMPORARY DESIGNATION OR EXPERIMENTAL NO.	
Western Plant Breeders, Inc	.	PH 986-61	WestBred 936
4. ADDRESS (street and no. or R.F.D. no., city, state, and ZIP) 8111 Timberline Drive		5. PHONE (include area code)	FOR OFFICIAL USE ONLY PVPO NUMBER
1.		(406) 587-1218	
Bozeman, Montana 59715			g Date
			FEBRUARY 28, 191
6. GENUS AND SPECIES NAME	Time A.M. D P.M.		
Triticum asetivum	Graminea	е	F Filing and Examination Fee:
8. CROP KIND NAME (Common Name)		9. DATE OF DETERMINATION	1 2, 32 <i>5.00</i>
Common Wheat (hard red spri	ng wheat)	May 1, 1989	Date
10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FOR	U DF ORGANIZATION	(Corporation, partnership,	Feb. 28, 1995
Corporation		•	Condition Fee:
11. IF INCORPORATED, GIVE STATE OF INCORPORATION		12. DATE OF INCORPORATION	v. 300.
Arizona		August 24, 199	0 July 19 1995
13. NAME AND ADORESS OF APPLICANT REPRESENTATIVE(S).	F ANY, TO SERVE I	THIS APPLICATION AND RECEIVE	ALL PAPERS
14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBLE A SEA Exhibit A, Origin and Breeding History of the Variety b. SOX Exhibit B, Novelty Statement c. XIX Exhibit C, Objective Description of Variety d. XIX Exhibit D, Additional Description of Variety e. XIX Exhibit E, Statement of the Basis of Applicant's Own f. XIX Sood Sample (2,500 viable untreated soods). Date g. XIX. Filing and Examination Fee (\$2,325) made payable	nership Seed Sample malled I to "Tressurer of the U	DO Plant Variety Protection Office	
15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VAR Plant Variety Protection Act) YES (# "YES," answer It	ETY BE SOLD BY VA ems 16 and 17 below	RIETY NAME ONLY AS A CLASS OF	CERTIFIED SEED? (See section 83(s) of the
16. DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE LIMITED AS TO NUMBER OF GENERATIONS? YES NO		ES" TO ITEM 16, WHICH CLASSES	OF PRODUCTION BEYOND BREEDER SEED?
18. DO THE APPLICANT(S) PREVIOUSLY FILE FOR PROTECTION	OF THE VARIETY IN	THE U.S.?	ISTERED CERTIFIED
□ Plant Variety Protection	r Act 📋 Pater	il Aci. Give date:	<u> </u>
18. HAS THE VARIETY BEEN RELEASED, USED, OFFERED FOR S YES (II "YES," GIVE NAMES OF COUNTRIES AND DAT	ESI USA M	ow the U.S. on other countries arch 15,1994	হ্ব
 The applicant(s) declare(s) that a viable sample of basic seeds such regulations as may be applicable. 	of this variety will be	turnished with the application and wil	Il be replonished upon request in accordance with
The undersigned applicant(s) is (are) the owner(s) of this sexuall in section 41, and is entitled to protection under the provisions	y reproduced novel pl of section 42 of the P	ant variety, and believe(s) that the valiant Variety Protection Act.	eriety is distinct, uniform, and stable as required
Applicant(s) is (are) informed that false representation herein ca			
SIGNATURE OF APPLICANT [OWNer(E)]		CAPACITY OR TITLE	DATE
(Vale R. Clark for Western Plant &	Breeders Inc.	Barley and Wheat Breeder	Feb. 24, 1995
SIGNATURE OF APPLICANT [OWING(S)]		APACITY OR TITLE	DATE
· Serving	1		
0.470			

14a. Origin and Breeding History

WestBred 936 is a hard red spring wheat selected by Western Plant Breeders from a male-sterile facilitated, recurrent selection population (MSFRSP) designated "906 alpha-84". This population was developed by top crossing WestBred 906-R onto male-sterile F2 plants in the two MSFRS populations designated SHRSPY-83 and EHRSP-83. These two populations were adapted primarily for the irrigated deserts of California and Arizona. Fifty such crosses were made near Phoenix, Arizona in March, 1984. The resultant F1 seed were bulked at harvest and planted at Bozeman, Montana in May, 1984. The F2 of this population was grown near Phoenix in the fall of 1984 and the spring of 1985. Selected spikes from this population were harvested, bulked, and planted as a bulk F3 near Bozeman in May, 1985. Individual spikes were harvested from this population in the fall of 1985 and planted as F4 rows near Phoenix in November, 1985. Agronomically acceptable rows were selected in May, 1986. One such row was given the experimental number PH 986-61. Seed from this row were planted near Bozeman in May, 1986 for further purification and increase. The resultant F5 plot (F6 seed) was harvested and used for yield testing in Arizona and California in 1987. Successive generations were tested in 1988, 1989, and 1990. PH 986-61 was too tall for these conditions and was dropped from further development for this area. PH 986-61 was yield tested in Idaho, Washington, and Montana in 1990, 1991, and 1992 and was found to be well adapted to these areas.

Spikes were selected near Phoenix in May, 1989 and were grown as head rows near Bozeman in the summer of 1989. Uniform rows were individually harvested and planted as large plots near Phoenix in the fall of 1989. Uniform plots were individually harvested in May, 1990. Seed from these plots was planted in individual strips near Bozeman in May, 1991. Uniform strips were harvested and the resultant seed was bulked to form Breeders seed. This Breeders seed was used to plant

WestBred 936

approximately 60 acres near Bozeman in the spring of 1992 for the purpose of producing Foundation seed. The production from this field was harvested in August, 1992 and designated "WestBred 936". Registered seed was planted in the spring of 1993 and certified seed was first sold to the general public March 15, 1994.

WestBred 936 is a stable and uniform variety in agronomic appearance and performance across several generations and growing conditions. Agronomic data to support this stability are presented in Tables 1 through 4.

VARIANTS:

A TACK VARIANT (one to Two head longth taller than the norm) occurs in West Bred 936 at a frequency of 3 per 10,000 plants. Also a white seed variant occurs at a frequency of up to 18 per 10,000 seeds.

14b. Novelty Statement

ман го Jone 1995 per letter

WestBred 936 is most similar to the variety WestBred 906-R. However, glume beak awn of WestBred 936 is twice the length of that of WestBred 906-R. Also, the kernels of WestBred 936 have a long brush where WestBred 906-R have a short brush, and WestBred 936 has a lax head where WestBred 906-R has a dense head. WestBred 936 is susceptible to the prevalent races of Hessian fly in the Pacific Northwest states of Washington and Idaho where WestBred 906-R is resistant. Another distinguishing character is that WestBred 936 is tolerant to the wild oat herbicide, difenzoquat ("Avenge") where WestBred 906-R is susceptible to damage. WestBred 936 is also approximately 4 cm. shorter than WestBred 906-R (t = 3.13 w21 df, p < .01).

The above comparisons, along with the complete objective description (14 c.), show WestBred 936 to be a novel variety of hard red spring wheat.

14c. Objective Description (see pages 4 and 5)

FORM APPROVED: OMB NO.0581-0055

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK AND SEED DIVISION BELTSVILLE, MARYLAND 20705

EXHIBIT C

OBJECTIVE DESCRIPTION OF VARIETY

	RITICUM SPP.)
MOSTON Dient Description	FOR OFFICIAL USE ONLY
Western Plant Breeders, Inc. ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	9500095
8111 Timberline Drive	VARIETY NAME OR TEMPORARY DESIGNATION
Bozeman, MT 59715	Wood-Drood 026
Place the appropriate number that describes the varietal charact	5/ -1 218 1
Place a zero in first box (e.g. 0 8 9 or 0 9) when numbe	r is either 99 or less or 9 or less.
I. KIND:	
1 1 = COMMON 2 = DURUM 3 = EMMER 4 = SPELT	5 = POLISH 6 = POULARD 7 = CLUB
2. TYPE,	
1 1 = SPRING 2 = WINTER 3 = OTHER (Specify)	1 = SOFT 3 = OTHER (Specify) 2 = HARD
2 1 = WHITE 2 = RED 3 = OTHER (Specify)	
3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO:	
0 5 2 FIRST FLOWERING	0 5 8 LAST FLOWERING
4. MATURITY (50% Flowering):	
0 6 NO. OF DAYS EARLIER THAN	. 7 1 = ARTHUR 2 = SCOUT 3 = CHRIS
0 2 NO. OF DAYS LATER THAN	. 8 4=LEMHI 5=NUGAINES 6=LEEDS 7 = Vandel 8= WestBred 906-R
PLANT HEIGHT (From soil level to top of head):	
0 8, 3 _{CM. HIGH}	
1 1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
1 5 CM. TALLER THAN	. [7]
0 5 CM. SHORTER THAN	1 = ARTHUR 2 = SCOUT 3 = CHRIS 7 = YecoraRc 4 = LEMHI 5 = NUGAINES 6 = LEEDS 8 = WestBred
PLANT COLOR AT BOOTING (See reverse):	4 = LEMHI 5 = NUGAINES 0 - ELESS 8 = WESTBYEC
3 1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN	1 1 = YELLOW 2 = PURPLE
STEM:	
1 Anthocyanin: 1 = ABSENT 2 = PRESENT	2 Waxy bloom: 1 = ABSENT 2 = PRESENT
2 Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT	1 Internodes: 1 = HOLLOW 2 = SOLID
0 4 NO. OF NODES (Originating from node above ground)	2 7 CM. INTERNODE LENGTH BETWEEN FLAG LEAF
AURICLES:	
2 Anthocyanin: 1 = ABSENT 2 = PRESENT	2 Hairiness: 1 = ABSENT 2 = PRESENT
LEAF:	
Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify):	2 Flag leaf: 1 = NOT TWISTED 2 = TWISTED
Hairs of first leaf sheath: = ABSENT 2 = PRESENT	2 Waxy bloom of flag leaf sheath: 1 = ABSENT 2 = PRESENT
1 9 MM. LEAF WIDTH (First leaf below flag leaf)	2 5 CM. LEAF LENGTH (First loaf below flag loaf):

06

11. HEAD:	· W		
1 Density, 1 = LAX	2 = DENSE	1 2 1	ING 2 = STRAP 3 = CLAVATE
4 Awnedness: 1 = AW	NLESS 2 = APICALLY AWNLETED	3 = AWNLETED 4 = AWNE	D
Color at maturity: 5	= WHITE 2 = YELLOW 3 = PINK 4 = BROWN 6 = BLACK 7 = OTHE	= RED R (Specify):	
1 1 CM. LENGTH		1 2 MM. WIDTH	
12. GLUMES AT MATURI	TV.		
3 Length: 1 = SHORT 3 = LONG((CA. 7 mm.) 2 = MEDIUM (CA. 8 mm.)	Width: 1 = NARROY 3 = WIDE (C	
1 7 1	ING 2 = OBLIQUE 3 = ROUNDED RE 5 = ELEVATED 6 = APICULATE	Beak: 1 = OBTUSE	2 = ACUTE 3 = ACUMINATE
13. COLEOPTILE COLOR		14. SEEDLING ANTHOCY	ANIN:
1 1 = WHITE 2 = R	ED 3 = PURPLE	1 = ABSENT 2	= PRESENT
15. JUVENILE PLANT GR	OWTH HABIT:	— I	
3 1 = PROSTRATE	2 = SEMI-ERECT 3 = EREC	э т ,	
16. SEED:			
1 Shape: l = OVATE	2 = OVAL 3 = ELLIPTICAL	1 Cheek: 1 = ROUND	ED 2 = ANGULAR
3 Brush: 1 = SHORT	2 = MEDIUM 3 = LONG	Brush: 1 = NOT CO	DLLARED 2 = COLLARED
O Phenol reaction (See instructions):	1 = IVORY 2 = FAWN 3 = LT. BROWN 4 = BROWN 5 = BLACK 0= Not		
Color: 1 = WHITE	2 = AMBER 3 = RED 4 = PURPLE		
0 7 MM. LENGTH	0. 3 MM. WIDTH	4 7 GM. PER 1000	SEEDS
17. SEED CREASE:			
2 Width: 1 = 60% OR L	ESS OF KERNEL 'WINOKA'	1 Depth: 1 = 20% Of	LESS OF KERNEL 'SCOUT'
1 .	ESS OF KERNEL 'CHRIS'	2 = 353 OF	LESS OF KERNEL 'CHRIS'
	AS WIDE AS KERNEL 'LEMHI'		LESS OF KERNEL 'LEMHI'
	·		
· ·	ted, 1 = Susceptible, 2 = Resistant)	·	
2 STEM RUST Preval	ent 2 (Races) Prevalent	2 stripe rust (Races) Preva	Lent 0 Loose SMUT
1 POWDERY MILDEW	0 BUNT	O OTHER (Specify)	
19. INSECT: (0 = Not Teste	d, 1 = Susceptible, 2 = Resistant)		
1 SAWFLY	O APHID (Bydv.)	0 GREEN BUG	O CEREAL LEAF BEETLE
OTHER (Specify)	· · · · · · · · · · · · · · · · · · ·	0 GP 0 A	0 B 0 c
Prevalent i Northwest	in the Pacific RACES: (0 D 0 E	0 F 0 G
20. INDICATE WHICH VARI	ETY MOST CLOSELY RESEMBLES THAT S	HBMITTED.	
CHARACTER	NAME OF VARIETY	CHARACTER	NAME OF VARIETY
Plant tillering		Seed size	
	WestBred 906-R		WestBred 906-R
Leaf size	11 11	Seed shape	71
	" "	Coleoptile elongation	
Leaf carriage	"	Seedling pigmentation	TI II

INSTRUCTIONS

GENERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form:

- (a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Whent Varieties Grown in the United States, Technical Bulletin 1278, United States Department of Agriculture.
- (b) W.E. Walls, 1965, A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysts. (See attachment.)
- LEAF COLOR: Nickerson's or any recognized color fan should be used to determine the leaf color of the described variety.

14d. Additional Description

WestBred 936 is a semi-dwarf, white chaffed, hard red spring wheat. The leaves and stems of WestBred 936 have a waxy bloom and the leaf auricles are purple with very few pubescent hairs. The spikes are lax, oblong and awned, and the awns are white at maturity. The glumes are white, long and wide, with narrow elevated shoulders. The beak is narrow, very long, and the apex is acuminate. Seed of WestBred 936 is mid-long, mid-wide, and ovate with rounded cheeks. The crease is mid-wide and shallow and the germ is medium in size. The brush is long and not collared. WestBred 936 is 1 to 2 inches shorter than WestBred 906-R and is less susceptible to lodging. WestBred 936 is more susceptible to powdery mildew (*Erysiphe graminis* f. sp. *tritici*) and head scab (*Fusarium* spp.) than WestBred 906-R. In contrast to W.B. 906-R, WestBred 936 appears to be tolerant to the wild oat herbicide "Avenge" (difenzoquat) in field tests.

Variants:

A tall variant (one to two head lengths taller than the norm) occurs in WestBred 936 at the frequency of 3 per 10,000 plants.

Also, a white seed variant occurs at a frequency of up to 18 per 10,000 seeds.

Western Plant Breeders is attempting to remove these variants through further head-rowing and purification. However, this may not be possible due to inherent genetic imbalances.

14e. Statement of ownership Replaced per totter MAH 20 June, 1995

Western Plant Breeders, Inc. is the employer of the breeder and is rightfully entitled to ownership and all rights of the variety "WestBred 936".

14e. Statement of ownership

'WestBred 936', the variety for which Plant Variety Protection is hereby sought, was developed by Dr. Dale R. Clark, an employee of Western Plant Breeders, Inc.. All rights to any invention, discovery, or development made by the employee while employed by Western Plant Breeders, Inc. were assigned by Western Plant Breeders, Inc. with no rights of any kind pertaining to 'WestBred 936' are retained by the employees

Table 1. Agronomic comparisons of WestBred 936 and check varieties in Western Plant Breeders trials.

Five Year Comparison

					4	•			
		Plt.Ht. <u>inches</u>	T.W. <u>lbs/bu</u>	Protein <u>%</u>	<u>BZ</u>	<u>BL</u>	<u>BU</u>	<u>ML</u>	4 Location <u>AVG</u>
W.B. 936	'91	34	60	14.2	8211	7509	6922	7093	7434
	'92	30	57	14.6	5554	6950	6867	6049	6355
	'93	33	61	12.7	6526	8078	6070	7014	6922
	'94	<u>29</u>	<u>62</u>	<u>14.2</u>	6994	<u>7985</u>	<u>6605</u>	6383	7121
	mean	32	60	13.9	6821	7631	6616	6635	6958
W.B. 906-R	'91	37	59	15.1	7795	6093	6486	6248	6655
	'92	30	57	14.4	5496	5943	5750	5631	5705
	'93	37	60	12.8	6867	6178	5804	7362	6553
	'94	<u>31</u>	<u>62</u>	14.4	<u>6704</u>	6693	<u>5973</u>	<u>5495</u>	<u>6297</u>
	mean	34	60	14.2	6716	6227	6003	6184	6302
W.B. Express	' 91	33	60	14.7	7338	6500	CC+D	0004	0500
Tribi Expicos	'92	29	58	14.7	7336 5902	6502 6814	6613 6104	6864 6137	6592
	'93	33	60	12.8	6445	7550	5398	7354	6239
	'94	<u>31</u>	<u>62</u>	13.9	<u>7158</u>	7654	6025	6724	6687 <u>7179</u>
	mean	32	60	14.0	6711	7130	6035	6770	6674
Fergus	· '91	37	60	14.1	8160	7200	6120	6780	6540
•	'92	32	59	14.1	5844	6814	6363	6382	6351
	'93	38	60	12.5	6990	8078	5085	7230	6846
	'94	<u>34</u>	<u>62</u>	14.2	<u>6720</u>	<u>7800</u>	6104	6250 6250	6711
	mean	35	60	13.7	6929	7473	5918	6661	6612

Table 2. Plant height in inches of WestBred 936 compared to WestBred 906-R in Western Plant Breeders' and public trials from 1990 - 1994.

<u>WPB</u>	WestBred 936	WestBred 906-R
1990 Bozeman,MT	29	32
Blackfoot,ID	29	30
Burley,ID	32	33
Moses Lake,WA 1991 Bozeman,MT Blackfoot,ID	38 36 31	39 39
Burley,ID Moses Lake,WA	34 36	32 33 40
Walla Walla,WA	35	36
1992 Bozeman,MT	30	31
Burley,ID	29	28
Moses Lake,WA	31	32
1993 Bozeman,MT	35	40
Burley,ID	31	34
Moses Lake,WA	34	37
1994 Bozeman,MT	29	31
Blackfoot,ID	34	34
Burley,ID	32	35
Moses Lake,WA	29	33
Univ. ID	00	24
1991 Aberdeen,ID	30	34
Tetonia,ID	28	28
CO St. Univ 1991 Center,CO	<u>37</u>	<u>38</u>
mean	32.68	34.05

t = 3.13 w/21 df, p < .01

Table 3. Agronomic comparisons of WestBred 936 and check varieties in Montana State Univ. trials from 1992 - 1994.

(25 location summary)

	l																
	∫awa	ď	8 8	3 8	3 6	8 8	3 &	- G	8 6	3 8	7 6		3 8	8	2	67	6
Yield (bu/ac)	21	r,	3 4) ú	8 6	, r	3 6	, rç	3 2	2 2	o N	27	, L	55	27	20	52
	83	7		3 6	5 8	8. 5	2 6	7	5 6	, <u>1</u>	8	9 6	8	67	7	9	8
	88	84	2	2 2	2	82	3 8	2 8	2 &	1 2	2	8 &	20.00	98	83	62	23
	awa	14.4	0.5	14.4	4.3	6	14.3	14.3	14.0	14.6		140	140	14.4	14.0	13.3	14.2
Protein %	54	6	14.9	5.	14.6	13.6	15.0	4 8	6.4	15.0	14.0	15.0	15.0	14.9	14.6	14.2	15.0
	83	5	13.8	14.4	13.9	12.5	4.4	14.0	13.7	14.7	13.1	13.7	13.4	14.8	4.2	13.1	13.9
	8	14.0	13.1	13.6		12.6	13.7	13.9	13.3	14.0	12.8	333	3.6	13.5	13.2	12.4	13.8
	gwg	58	23	28	28	23	8	8	8	28	28	26	28	23	23	29	29
T.W. (lbs/bu)	94	28	26	23	26	29	8	8	20	29	28	23	28	29	26	29	20
	88	26	28	27	22	57	8	200	26	22	22	28	22	28	28	28	28
	85	9	9	9		8	9	9	6	8	26	6	8	09	8	9	61
6	avg	59	હ	စ္က	88	႘	89	89	ဗ္ဗ	ဗ္ဗ	g	ဓ	34	88	37	32	32
Pit. Ht. (inche:	94	28	99	8	27	ဓ	37	37	8	ਲ	ਲ	53	32	37	98	8	હ
	63	30	8	စ္က	58	ဗ္ဗ	99	æ	ස	34	34	હ	33	88	æ	8	ဗ္ဗ
,	8	59	32	ဓ		32	96 96	æ	ဗ္ဗ	ဗ္ဗ	ဗ္ဗ	ဓ	34	ဗ္ဗ	စ္တ	32	8
	SVG	168	169	168	173	174	175	172	7	173	173	2	173	172	172	173	168
Date 1/1	84	166	167	166	168	173	173	169	168	170	171	167	171	169	170	171	166
Heading Date from 1/1	ଷ	172	175	172	178	2	180	177	176	178	178	175	177	177	177	179	174
	8	166	165	165		7	171	169	168	169	120	166	13	169	169	169	164
Variety		WB 936	Fergus	WB 926	WB Express	Newana	Lew	Fortuna	Pondera	Len	Glenman	Hi-Line	McNeal	Stoa	Amidon	Rambo	Border

Table 4. Milling and baking quality data summary of WestBred 936 compared to check varieties in Montana State University's Intrastate Wheat Yield Trials in 1992 and 1993.

	G & T		6.2	6.1	5.4	го С	9.53	5.7			. 1.9	5.9	5.8	5.9	6.1	5.6
<u> 120</u>	LF VL		1004	1019	949	955	1003	1008	·		935	931	937	891	941	872
Baking Data	MIX		2.8	2.9	2.3	23	3.5	3.4			4.0	6 6	3.4	3.0	4.9	4.6
- 1	ABS%		66.7	66.4	67.1	9.99	68.4	2'69			65.2	65.4	2'99	65.6	67.5	67.8
	E E		14	Ξ	22	56	165	66			20	8	19	25	र	ω
- <u>-</u> -	SIAB		10.0	12.5	6.3	6.3	9.3	11.4			11.9	11.1	8.8	7.4	10.7	12.0
Farinograph	ZEAK		9.1	6.7	5.6	5.6	7.3	10.9		•	10.9	10.9	6.4	5.8	5.9	7.2
	ABS %		66.3	65.6	67.1	64.7	67.8	69.3			63.2	63.9	65.3	63.4	65.3	66.3
Flour	A30 %		0,460	0.453	0.475	0.424	0.446	0.461			0.429	0.476	0.473	0.445	0.438	0.458
9	1 LU %		71.6			68.3	69.1	69.5			68.3	6.69	8.69	9.79	66.7	68.6
		1992 (7 location avg)	WestBred 936	WestBred 926	Fergus	Newana	Hi-Line	McNeal		1993 (7 location avg)	WestBred 936	WestBred 926	Fergus	Newana	Hi-Line	McNeal